



## Partial Characterization of $\alpha$ -Galactosidic Activity from the Antarctic Bacterial Isolate, *Paenibacillus* sp. LX-20 as a Potential Feed Enzyme Source

Inkyung Park, Jaekoo Lee and Jaiesoon Cho\*

Department of Animal Sciences and Environment, College of Animal Bioscience and Technology,  
Konkuk University, Seoul 143-701, Korea

**ABSTRACT:** An Antarctic bacterial isolate displaying extracellular  $\alpha$ -galactosidic activity was named *Paenibacillus* sp. LX-20 based on 16S rRNA gene sequence analysis. Optimal activity for the LX-20  $\alpha$ -galactosidase occurred at pH 6.0-6.5 and 45°C. The enzyme immobilized on the smart polymer Eudragit L-100 retained 70% of its original activity after incubation for 30 min at 50°C, while the free enzyme retained 58% of activity. The enzyme had relatively high specificity for  $\alpha$ -D-galactosides such as *p*-nitrophenyl- $\alpha$ -galactopyranoside, melibiose, raffinose and stachyose, and was resistant to some proteases such as trypsin, pancreatin and pronase. Enzyme activity was almost completely inhibited by Ag<sup>+</sup>, Hg<sup>2+</sup>, Cu<sup>2+</sup>, and sodium dodecyl sulfate, but activity was not affected by  $\beta$ -mercaptoethanol or EDTA. LX-20  $\alpha$ -galactosidase may be potentially useful as an additive for soybean processing in the feed industry. (**Key Words:** Antarctic, *Paenibacillus* sp.  $\alpha$ -Galactosidase, Feed Industry)

### INTRODUCTION

Soybean meal (SBM) is the main protein supplement for poultry and livestock feed worldwide (Ghazi et al., 2003; Stein et al., 2008) and comprises over 50% of the global protein sources in poultry and livestock feed (Kohlmeier, 1990). Moreover, SBM galactooligosaccharides have contributed to improving intestinal health of animals by increasing concentrations of beneficial gut bacteria including bifidobacteria and lactobacilli, as well as concentrations of short-chain fatty acids, which might be a potential alternative for subtherapeutic antibiotic supplementation (Smiricky-Tjardes et al., 2003). However, SBM  $\alpha$ -galactooligosaccharides such as raffinose (galactose- $\alpha$ -1,6-sucrose), and stachyose (galactose- $\alpha$ -1,6-raffinose) which constitute about 1% and 6% on dry matter basis, respectively (Grieshop et al., 2003), act as anti-nutritive factors that may depress animal performance (Anderson and Wolf, 1995). The oligosaccharides appear to increase viscosity of digesta, hindering the digestion of nutrients by reducing their interaction with digestive

enzymes in the small intestine (Smits and Annison, 1996). Since monogastric animals including humans have very limited inherent  $\alpha$ -galactosidase (EC 3.2.1.22,  $\alpha$ -D-galactoside galactohydrolase), which is essential to hydrolyze  $\alpha$ -galactosidic linkages in the aforementioned sugars, they are transferred intact into the large intestine, where anaerobic bacteria ferment them and lead to intestinal disturbances such as flatulence (Falkoski et al., 2006; Yoon and Hwang, 2008). Therefore, the reduction of these oligosaccharides in SBM by microbial  $\alpha$ -galactosidase treatment can potentially improve the nutritive value of SBM and likely helps to improve the production performance of the animals fed SBM-containing diets (Ghazi et al., 2003; Falkoski et al., 2006; Ao et al., 2009).

Microorganisms are the most attractive sources for industrial enzyme production (Syed et al., 2009). Microbially-enhanced feed production has greatly benefited from biotechnological approaches (Hasan et al., 2006). Many microbial  $\alpha$ -galactosidases have been extensively studied regarding their enzymatic properties and the genes encoding  $\alpha$ -galactosidases have been cloned (Fridjonsson et al., 1999; Yoon and Hwang, 2008). However, little is known of the  $\alpha$ -galactosidic activity of *Paenibacillus* sp., which has been regarded as an attractive source for producing

\* Corresponding Author: Jaiesoon Cho. Tel: +82-2-450-3375, Fax: +82-2-455-1044, E-mail: chojs70@konkuk.ac.kr  
Submitted Dec. 23, 2011; Accepted Feb. 17, 2012; Revised Mar. 5, 2012

carbohydrate degrading enzymes such as cellulase, xylanase, and  $\beta$ -glucosidase (Shipkowski and Brenchley, 2005; Kim et al., 2009; Waeonukul et al., 2009). The purpose of this study was to explore the partial catalytic properties of  $\alpha$ -galactosidase activity from an Antarctic bacterial isolate, *Paenibacillus sp.* LX-20, as a potential feed enzyme source.

## MATERIALS AND METHODS

### Bacterial strain and culture conditions

LX-20 derived from Antarctic soil samples was supplied by the Korea Polar Research Institute operating the King Sejong Station (South Korea) in Antarctica. Screening for  $\alpha$ -galactosidase activity was performed by the appearance of blue colonies on selective Luria Bertani (LB) agar (pH 7.2) supplemented with 0.2% lactose and 32  $\mu$ g/ml of 5-bromo-4-chloro-3-indolyl- $\alpha$ -D-galactopyranoside (X- $\alpha$ -Gal) as previously described (Goulas et al., 2009) at 28°C. Growth and  $\alpha$ -galactosidase production were investigated in an Erlenmeyer flask (2 L) containing 500 ml of the enzyme production medium composed of Bacto tryptic soy broth (Difco) (1.7% pancreatic digest of casein, 0.3% papaic digest of soybean, 0.25% dextrose, 0.5% NaCl, 0.25% dipotassium phosphate; pH 7.2) and 0.3% soybean meal inoculated with 1% (vol/vol) 24 h inoculum at 28°C with vigorous shaking (220 rpm) by monitoring the absorbance (O.D.<sub>600nm</sub>) and  $\alpha$ -galactosidase activity of the culture supernatant at various time points.

### Taxonomic identification of strain LX-20

Genomic DNA was extracted from strain LX-20 using the FastDNA kit (Qbiogene) according to the manufacturer's protocol. The 16S rRNA gene was amplified from genomic DNA by PCR using the universal primers, 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-GGTACCTTGTTACGACTT-3') (William et al., 1991). The amplified 1,453 bp sequences were determined by an automated ABI PRISM 3730 XL DNA analyzer (Applied Biosystems). The resulting sequences were compared with the GenBank database (NCBI) using BLAST (Altschul et al., 1990). Sequences showing a relevant degree of similarity were imported into the CLUSTAL W program (Thompson et al., 1994) and aligned. The evolutionary distances with other strains of *Paenibacilli* were computed using the Maximum Composite Likelihood method (Tamura et al., 2004) and the phylogenetic relationships were determined using the software MEGA version 4.0 (Tamura et al., 2007).

### Partial purification of the enzyme

Strain LX-20 was cultivated in 1 L of the enzyme production medium for 96 h at 28°C. The culture medium

containing secreted  $\alpha$ -galactosidase was centrifuged (10,000 $\times$ g; 30 min; 4°C) to remove cells, and the protein in the supernatant was then precipitated with ammonium sulfate (50% saturation). The pellet was dissolved in 50 mM Tris-HCl (pH 8.0) and dialyzed overnight against 50 mM Tris-HCl (pH 7.4) at 4°C. The dialyzed solution was used as the enzyme source to examine the catalytic properties throughout this work.

### Zymogram analysis

The enzyme was subjected to non-denaturing 6.5% polyacrylamide gel electrophoresis (PAGE) using a Modular Mini-Protein II Electrophoresis System (Bio-Rad) according to the manufacturer's instructions. After gel electrophoresis, the gel was placed on 1.5% (wt/vol) bacto agar plate containing 4 mg/ml X- $\alpha$ -Gal and was incubated at 40°C for 12 h. The band of  $\alpha$ -galactosidase activity was detected by appearance of a blue zone.

### Enzyme assay and substrate specificity

Unless otherwise stated,  $\alpha$ -galactosidase activity was measured at 40°C by assaying the release of *p*-nitrophenol from *p*-nitrophenyl- $\alpha$ -D-galactopyranoside with a final concentration of 1 mM in 1 ml of 50 mM sodium phosphate (pH 6.5). Activity on other *p*-nitrophenyl (*p*NP) or *o*-nitrophenyl (*o*NP) conjugated synthetic substrates such as *p*NP- $\alpha$ -L-arabinofuranoside, *p*NP- $\alpha$ -D-glucopyranoside, *p*NP- $\beta$ -D-galactopyranoside, *p*NP- $\beta$ -D-xylopyranoside, *p*NP- $\beta$ -D-cellobioside, *o*NP- $\beta$ -D-galactopyranoside, and *o*NP- $\beta$ -D-glucopyranoside was determined in 1 ml of 50 mM sodium phosphate (pH 6.5) at 40°C with a final substrate concentration of 1 mM. The reactions were then stopped by the addition of 1 ml of 1 M Na<sub>2</sub>CO<sub>3</sub> and color development was measured at O.D.<sub>405 nm</sub>. One unit (U) of enzyme activity was defined as the amount of enzyme to produce 1  $\mu$ mol of *p*-nitrophenol or *o*-nitrophenol per minute under the assay conditions described. When maltose, carboxymethylcellulose (CMC), xylan, galactomannan and starch were used as the substrate with a final concentration of 0.4%, the production of reducing sugar was determined under standard assay conditions by the dinitrosalicylic acid (DNS) method (Miller, 1959). Determination of activity against lactose, raffinose, melibiose, and stachyose was evaluated by assaying the release of D-galactose using a galactose test kit (Boehringer Mannheim GmbH). One enzyme unit (U) was defined as the amount of enzyme to produce 1  $\mu$ mol of reducing sugar equivalent, or galactose per minute under the assay conditions.

### Effect of pH and temperature on enzyme activity

$\alpha$ -Galactosidase activities were investigated in the pH range of 3-9 (50 mM glycine-HCl (pH 3); 50 mM sodium

acetate (pH 4-5.5); 50 mM sodium phosphate (pH 5.5-7); 50 mM Tris-HCl (pH 7-9)) at 30°C and temperatures between 0 and 80°C at the optimum pH.

#### Determination of protease resistance

The stability of LX-20  $\alpha$ -galactosidase to proteolysis was investigated as previously described (Cao et al., 2010) with slight modification. The partially purified enzyme was incubated with trypsin, pancreatin, proteinase K, subtilisin Carlsberg or pronase (all from Sigma) at 37°C for 30 min in 0.1 M sodium phosphate (pH 7.0), using a protease:  $\alpha$ -galactosidase (wt:wt) ratio of 1:10 and then subjected to enzyme activity assay under standard conditions.

#### Effect of metal ions and chemicals on enzyme activity

The effect of different metal ions and chemicals on  $\alpha$ -galactosidase activity was determined under standard assay conditions after the enzyme was pre-incubated in the presence of 1 mM of  $Mg^{2+}$ ,  $Ca^{2+}$ ,  $Co^{2+}$ ,  $Ni^{2+}$ ,  $Cu^{2+}$ ,  $Mn^{2+}$ ,  $Fe^{2+}$ ,  $Hg^{2+}$ ,  $Ba^{2+}$ ,  $Ag^+$ ,  $K^+$ ,  $Na^+$ ,  $\beta$ -mercaptoethanol, EDTA, SDS, or PMSF (phenylmethylsulfonyl fluoride) in 50 mM sodium phosphate (pH 6.5) for 20 min at 25°C.

#### Preparation of immobilized enzyme

2% (wt/vol) Eudragit L-100 (Rohm Pharma, Weiterstadt, Germany) solution was prepared as previously described (Roy et al., 2003). Partially purified LX-20  $\alpha$ -galactosidase (2.5 ml) was added to 0.75 ml of the Eudragit L-100 solution and the final volume was made up to 5.0 ml with 50 mM sodium phosphate (pH 6.5). After incubation for 1 h at 25°C, the polymer was precipitated by lowering the pH to 4.0 with 3 M acetic acid. After 20 min, the suspension was centrifuged at 12,000 $\times$ g for 20 min. The pellet was washed with 4 ml of 10 mM sodium acetate (pH 4.0) and the suspension was again centrifuged. Finally, the pellet was suspended in 4 ml of 50 mM sodium phosphate (pH 6.5) and used as the immobilized enzyme. As a reference, the stability of the enzyme at pH 4 during immobilization process was monitored.


#### Thermal stability

Thermal stability of the LX-20  $\alpha$ -galactosidase was estimated by incubation of the free enzyme or the immobilized enzyme in 50 mM sodium phosphate (pH 6.5) for 30 min at various temperatures ranging from 30 to 80°C before measuring.

## RESULTS AND DISCUSSION

#### Identification of isolated strain LX-20 and the $\alpha$ -galactosidase production

To identify the isolated strain LX-20 showing



**Figure 1.** Zymogram analysis of  $\alpha$ -galactosidase activity in the enzyme preparation.

$\alpha$ -galactosidase activity (Figure 1), the 16S rRNA gene was cloned and the sequence was compared with those available in the database. A phylogenetic tree based on the 16S rRNA gene sequences from 11 *Paenibacillus* strains showed that strain LX-20 shared 99.1% sequence identity with the type strain, *Paenibacillus odorifer* DSM 15391<sup>T</sup> (Figure 2). Therefore, it was named *Paenibacillus* sp. LX-20.


Extracellular  $\alpha$ -galactosidase production by strain LX-20 was somewhat growth phase-dependent (Figure 3). During log phase of growth, little  $\alpha$ -galactosidase was secreted, but about 30-fold increase in extracellular  $\alpha$ -galactosidase activity was induced during transition to stationary phase of growth, reaching a maximum (0.084 $\pm$ 0.00019 U/ml) at 36 h of incubation. Meanwhile, cell growth steeply increased after the initial 8 h of incubation and reached a peak (4.35 $\pm$ 0.03 of O.D<sub>600 nm</sub>) at 28 h of incubation.

#### Nucleotide sequence accession numbers

The nucleotide sequence of the 16S rRNA gene has been deposited in the GenBank database under Accession No. HQ660810.

#### Effect of pH on enzyme activity

As shown in Figure 4A, the optimal activity for the LX-20  $\alpha$ -galactosidase occurred at pH 6.0-6.5, and over 45% of the activity was achieved between pH 5.5 and 7.0. The enzyme was nearly inactivated at acidic pH ranges (3 to 5), which characteristic of most bacterial  $\alpha$ -galactosidases (Leder et al., 1999; King et al., 2002; Patil et al., 2010), even though fungal and yeast  $\alpha$ -galactosidases generally display optimal activities in acidic conditions (Viana et al., 2006; Wang et al., 2010). In fact, the acidic  $\alpha$ -galactosidase is inappropriate for the removal of galactooligosaccharides in soymilk, which can be used as an economic partial substitute for whole milk in calf-raising facilities (Ghorbani




**Figure 2.** Phylogenetic relationship of the 16S rRNA sequences of *Paenibacillus* sp. LX-20 with other type strains of *Paenibacillus*. Bootstrap values (based on 1,000 trials and only values >50%) are shown at the nodes. The GenBank accession numbers are indicated in parentheses. Bar denotes 5-base substitutions per 1,000 nucleotide position. *P.* is an abbreviation of *Paenibacillus*.

et al., 2007), because the natural pH of soymilk is 6.2 to 6.4 (Yoon and Hwang, 2008). Nevertheless,  $\alpha$ -galactosidase from a bacterial strain, *Leuconostoc mesenteroides* JK55 can retain 70% of its maximal activity at pH 4.0 (Yoon and Hwang, 2008).


#### Effect of temperature on enzyme activity and thermal stability

LX-20  $\alpha$ -galactosidase showed optimal activity at 45°C and more than 55% of the highest activity remained at 30

45°C (Figure 4B), which is reminiscent of enzymes from mesophilic microbes. Generally, enzymes produced by microbes that dwell in cold environments display higher catalytic efficiency at low temperatures and greater thermosensitivity than their mesophilic counterparts (Gerday et al., 1997). For example, CelG from the Antarctic bacterium, *Pseudoalteromonas haloplanktis*, is a heat-labile cellulase and its half-life during incubation at 45°C is approximately 40 min (Garsoux et al., 2004). Meanwhile, a cold-active cellulase, CelG can be produced from the



**Figure 3.** Growth and  $\alpha$ -galactosidase production by *Paenibacillus* sp. LX-20. Symbols represent  $\alpha$ -galactosidase activity (dotted line) and growth (solid line). Data were expressed as mean and standard errors from three experiments.




**Figure 4.** Optimal pH (A) and temperature (B) activity profiles. (A) Relative activity at 30°C and various pHs where 100% equates to  $0.032\pm 0.0011$  U/ml. Used buffers : 50 mM glycine-HCl (pH 3) (closed triangle), 50 mM sodium acetate (pH 4 to 5.5) (closed triangle), 50 mM sodium phosphate (pH 5.5 to 7) (closed square), 50 mM Tris-HCl (pH 7 to 9) (closed triangle) (B) Relative activity at pH 6.5 and various temperatures where 100% equates to  $0.035\pm 0.0003$  U/ml. The assays were performed at a final concentration of 1 mM *p*-Np- $\alpha$ -galactopyranoside. Data were expressed as mean and standard errors from three experiments.

mesophilic ruminal anaerobe, *Fibrobacter succinogenes* S85, showing a temperature optimum of 25°C and complete inactivation even after 20 min of exposure at 50°C (Iyo and Forsberg, 1999). In fact, LX-20  $\alpha$ -galactosidase may be suitable for the use of a feed supplement to poultry or swine diets because the optimal temperature range of enzyme is close to the intestinal temperature of the animals (37 to 40°C) (Lei and Porres, 2003). LX-20  $\alpha$ -galactosidase was successfully immobilized using Eudragit L-100 and the enzyme was stable at pH 4 during the immobilization process (Figure 5). Even if the immobilization of LX-20  $\alpha$ -galactosidase on the smart polymer Eudragit (Roy et al., 2003; Ai et al., 2005) had no remarkable effect on the thermal stability of the enzyme, the immobilized enzyme


retained 70% of its original activity after incubation for 30 min at 50°C, while the corresponding activity of the free enzyme was only 58% (Figure 6).

#### Effect of proteolysis on enzyme activity


LX-20  $\alpha$ -galactosidase showed different resistances to the tested proteases (Figure 7). The enzyme retained about 50% and 60% of its initial activity after 30 min incubation with proteinase K and subtilisin Carlsberg, respectively. However, trypsin, pancreatin and pronase had no effect on the enzymatic activity. The susceptibility of feed enzymes to proteolytic attack is important because the rate and site of inactivation of an enzyme are determined by this property (Wang et al., 2007). In contrast, the bacterial  $\alpha$ -galactosidase



**Figure 5.** The stability of the LX-20  $\alpha$ -galactosidase at pH 4 during immobilization process. Data were expressed as mean and standard errors from three experiments.



**Figure 6.** Temperature effects on the stability of the LX-20  $\alpha$ -galactosidase. Symbols represent the free form enzyme (dotted line) and the immobilized form enzyme (solid line). Data were expressed as mean and standard errors from three experiments.



**Figure 7.** Effect of proteases on the LX-20  $\alpha$ -galactosidase activity. The activity of protease-untreated control was defined as 100%, which equates to  $0.034 \pm 0.009$  U/ml. Data were expressed as mean and standard errors from three experiments.

from *Streptomyces sp. S27* loses about 50% of its initial activity after 30 min of trypsin digestion (Cao et al., 2010).

#### Effect of metal ions and inhibitors on enzyme activity

LX-20  $\alpha$ -galactosidase was almost completely inhibited by  $\text{Ag}^+$ ,  $\text{Hg}^{2+}$ ,  $\text{Cu}^{2+}$ , and SDS (Table 1). This is consistent with previous result reported for fungal  $\alpha$ -galactosidase from *Penicillium griseoroseum* (Falkoski et al., 2006). The LX-20  $\alpha$ -galactosidase appears to possess sensitive sulfhydryl groups in the active site, since  $\text{Ag}^+$  and  $\text{Hg}^{2+}$  are strong thiol-specific inhibitors of several  $\alpha$ -galactosidases (King et al., 2002; Cao et al., 2010; Wang et al., 2010). The hypersensitivity of the enzyme to the common anionic detergent SDS also indicates that hydrogen bonds may play a pivotal role in maintaining the enzymatic activity (Wang et al., 2005). However,  $\text{Cu}^{2+}$  stimulated the activity of  $\alpha$ -galactosidase from an acidophilic fungus, *Bispora sp.* (Wang et al., 2010).  $\text{Ca}^{2+}$ ,  $\text{Co}^{2+}$ ,  $\text{Mn}^{2+}$ ,  $\text{Ba}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$ , and  $\text{K}^+$  had no major effects on the activity of LX-20  $\alpha$ -galactosidase. The enzyme was not affected by  $\beta$ -mercaptoethanol and EDTA, suggesting that it may not be a metalloenzyme (Falkoski et al., 2006; Viana et al., 2006). Additionally, the enzyme was partially inhibited by  $\text{Fe}^{2+}$ , and  $\text{Ni}^{2+}$ , and the established serine protease inhibitor PMSF (Hutadilok-Towatana et al., 1999).

#### Substrate specificity

LX-20  $\alpha$ -galactosidase activity against various substrates was summarized in Table 2. The enzyme was specific for the aryl- $\alpha$ -galactosidic substrate, *p*-nitrophenyl-

**Table 1.** Effect of metal ions and chemicals on LX-20  $\alpha$ -galactosidase activity

Reagent <sup>a</sup>	Relative activity (%) <sup>b</sup>
No addition	100.0
$\text{Ca}^{2+}$	100.6±1.1
$\text{Co}^{2+}$	84.1±1.6
$\text{Fe}^{2+}$	74.3±1.1
$\text{Mg}^{2+}$	101.3±2.1
$\text{Mn}^{2+}$	85.8±2.3
$\text{Ba}^{2+}$	89.1±0.7
$\text{Cu}^{2+}$	1.9±1.8
$\text{Ni}^{2+}$	34.5±0.6
$\text{Hg}^{2+}$	17.0±4.0
$\text{K}^+$	96.6±0.2
$\text{Na}^+$	99.1±0.9
$\text{Ag}^+$	0.0
$\beta$ -mercaptoethanol	110.0±0.8
EDTA	102.3±0.3
SDS	2.7±1.2
PMSF	69.1±1.3

<sup>a</sup> The final concentration of each reagent was 1 mM in the assay buffer.

<sup>b</sup> 100% was assigned to the activity in the absence of all reagents and equates to 0.037±0.0003 U/ml. Data were expressed as mean and standard errors from three experiments.

$\alpha$ -galactopyranoside. However, little or no activity was detected against maltose, lactose and other synthetic substrates containing  $\beta$ -linkages or containing arabinose and glucose residues. The enzyme showed no ability to hydrolyze polysaccharides such as carboxymethylcellulose,

**Table 2.** Substrate specificity of LX-20  $\alpha$ -galactosidase

Substrates	Concentration	Enzyme activity (U/mL)
<i>p</i> -nitrophenyl- $\alpha$ -D-galactopyranoside	1 mM	0.033±0.001
<i>p</i> -nitrophenyl- $\beta$ -D-galactopyranoside	1 mM	0.002±0.0001
<i>p</i> -nitrophenyl- $\alpha$ -D-glucopyranoside	1 mM	0.000
<i>o</i> -nitrophenyl- $\beta$ -D-galactopyranoside	1 mM	0.000
<i>p</i> -nitrophenyl- $\beta$ -D-xylopyranoside	1 mM	0.000
<i>p</i> -nitrophenyl- $\beta$ -D-cellobioside	1 mM	0.000
<i>o</i> -nitrophenyl- $\beta$ -D-glucopyranoside	1 mM	0.000
<i>p</i> -nitrophenyl- $\alpha$ -L-arabinofuranoside	1 mM	0.004±0.002
Raffinose	1 mM	0.752±0.024
Melibiose	1 mM	2.476±0.073
Stachyose	1 mM	2.816±0.088
Lactose	1 mM	0.000
Carboxymethylcellulose (CMC)	0.4%	0.000
Xylan (birchwood)	0.4%	0.000
Galactomannan (locust bean gum)	0.4%	0.000
Starch	0.4%	0.000
Maltose	0.4%	0.000

Data were expressed as mean and standard errors from three experiments.

xylan, galactomannan (locust bean gum) and starch. Among the galacto-oligosaccharides tested, stachyose was most effectively hydrolyzed by the enzyme in comparison with raffinose or melibiose. In contrast, most  $\alpha$ -galactosidases degrade raffinose rapidly and stachyose slowly (Ishiguro et al., 2001). Many family 27  $\alpha$ -galactosidases catalyze the release of galactose from intact galactomannan polymers, whereas the substrate specificity in family 36 is restricted to small oligosaccharides including raffinose and stachyose (Wang et al., 2010). Thus, the LX-20  $\alpha$ -galactosidase may belong to family 36.

In conclusion, LX-20  $\alpha$ -galactosidase may be potential for use as an additive for soybean processing in the feed industry, because the temperature and pH optima of the enzymatic activity are appropriate for the physiological surroundings and manufacturing conditions in soybean processing, and the enzyme showed high specificity for galacto-oligosaccharides, as well as resistance to some intestinal proteases. Future scientific studies including gene cloning, protein engineering, and downstream fermentation technology will focus on maximizing the catalytic efficiency and productive yield of the enzyme.

#### ACKNOWLEDGEMENT

This paper was supported by Konkuk University in 2011.

#### REFERENCES

- Ai, Z., Z. Jiang, L. Li, W. Deng, I. Kusakabe and H. Li. 2005. Immobilization of *Streptomyces olivaceoviridis* E-86 xylanase on Eudragit S-100 for xylo-oligosaccharide production. *Process Biochem.* 40:2707-2714.
- Altschul, S. F., W. Gish, W. Miller, E. W. Myers and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403-410.
- Anderson, R. L. and W. J. Wolf. 1995. Compositional changes in trypsin inhibitors, phytic acid, saponins and isoflavones related to soybean processing. *J. Nutr.* 125:581S-588S.
- Ao, T., A. H. Cantor, A. J. Pescatore, M. J. Ford, J. L. Pierce and K. A. Dawson. 2009. Effect of enzyme supplementation and acidification of diets on nutrient digestibility and growth performance of broiler chicks. *Poult. Sci.* 88:111-117.
- Cao, Y., T. Yuan, P. Shi, H. Luo, N. Li, K. Meng, Y. Bai, P. Yang, Z. Zhou, Z. Zhang and B. Yao. 2010. Properties of a novel  $\alpha$ -galactosidase from *Streptomyces sp.* S27 and its potential for soybean processing. *Enzyme Microb. Technol.* 47:305-312.
- Falkowski, D. L., V. M. Guimaraes, C. M. Callegari, A. P. Reis, E. G. De Barros and S. T. De Rezende. 2006. Processing of soybean products by semipurified plant and microbial  $\alpha$ -galactosidases. *J. Agric. Food Chem.* 54:10184-10190.
- Fridjonsson, O., H. Watzlawick, A. Gehweiler, T. Rohrhirsch and R. Mattes. 1999. Cloning of the gene encoding a novel thermostable  $\alpha$ -galactosidase from *Thermus brockianus* ITI360. *Appl. Environ. Microbiol.* 65:3955-3963.
- Garsoux, G., J. Lamotte, C. Gerday and G. Feller. 2004. Kinetic and structural optimization to catalysis at low temperatures in a psychrophilic cellulase from the Antarctic bacterium *Pseudoalteromonas haloplanktis*. *Biochem. J.* 384:247-253.
- Gerday, C., M. Aittaleb, J. L. Arpigny, E. Baise, J. P. Chessa, G. Garsoux, I. Petrescu and G. Feller. 1997. Psychrophilic enzymes: a thermodynamic challenge. *Biochim. Biophys. Acta* 1342:119-131.
- Ghazi, S., J. A. Rooke and H. Galbraith. 2003. Improvement of the nutritive value of soybean meal by protease and  $\alpha$ -galactosidase treatment in broiler cockerels and broiler chicks. *Br. Poult. Sci.* 44:410-418.
- Ghorbani, G. R., R. Kowsar, M. Alikhani and A. Nikkhah. 2007. Soymilk as a novel milk replacer to stimulate early calf starter intake and reduce weaning age and costs. *J. Dairy Sci.* 90:5692-5697.
- Goulas, T., A. Goulas, G. Tzortzis and G. R. Gibson. 2009. A novel  $\alpha$ -galactosidase from *Bifidobacterium bifidum* with transgalactosylating properties: gene molecular cloning and heterologous expression. *Appl. Microbiol. Biotechnol.* 82:471-477.
- Griehop, C. M., C. T. Kadzere, G. M. Clapper, E. A. Flickinger, L. L. Bauer, R. L. Frazier and G. C. Fahey Jr. 2003. Chemical and nutritional characteristics of United States soybeans and soybean meals. *J. Agric. Food Chem.* 51:7684-7691.
- Hasan, F., A. A. Shah and A. Hameed. 2006. Industrial applications of microbial lipases. *Enzyme Microb. Technol.* 39:235-251.
- Hutadilok-Towatana, N., A. Painupong and P. Suintanalert. 1999. Purification and characterization of an extracellular protease from alkaliphilic and thermophilic *Bacillus sp.* PS 719. *J. Biosci. Bioeng.* 87:581-587.
- Ishiguro, M., S. Kaneko, A. Kuno, Y. Koyama, S. Yoshida, G. G. Park, Y. Sakakibara, I. Kusakabe and H. Kobayashi. 2001. Purification and characterization of the recombinant *Thermus sp.* T2  $\alpha$ -galactosidase expressed in *Escherichia coli*. *Appl. Environ. Microbiol.* 67:1601-1606.
- Iyo, A. H. and C. W. Forsberg. 1999. A cold-active glucanase from the ruminal bacterium *Fibrobacter succinogenes* S85. *Appl. Environ. Microbiol.* 65:995-998.
- Kim, B. C., K. H. Lee, M. N. Kim, E. M. Kim, S. R. Min, H. S. Kim and K. S. Shin. 2009. *Paenibacillus pini sp. nov.*, a cellulolytic bacterium isolated from the rhizosphere of pine tree. *J. Microbiol.* 47:699-704.
- King, M. R., B. A. White, H. P. Blaschek, B. M. Chassy, R. I. Mackie and I. K. O. Cann. 2002. Purification and characterization of a thermostable  $\alpha$ -galactosidase from *Thermoanaerobacterium polysaccharolyticum*. *J. Agric. Food Chem.* 50:5676-5682.
- Kohlmeier, R. H. 1990. World production, storage and utilization of various defatted animal and vegetable mid-high protein meals. In: *World Conference of Edible Fats Oils Processing Basic Principles and Modern practices* (Ed. D. R. Erikson). Am. Oil Chem. Soc. Champaign, IL, USA. p. 390.
- Leder, S., W. Hartmeier and S. P. Marx. 1999.  $\alpha$ -galactosidase of *Bifidobacterium adolescentis* DSM 20083. *Curr. Microbiol.* 38:101-106.
- Lei, X. G. and J. M. Porres. 2003. Phytase enzymology,

- applications and biotechnology. *Biotechnol. Lett.* 25:1787-1794.
- Miller, G. L. 1959. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.* 31:426-428.
- Patil, A. G. G., S. K. Praveen Kumar, V. H. Mulimani, Y. Veeranagouda and K. Lee. 2010.  $\alpha$ -galactosidase from *Bacillus megaterium* VHM1 and its application in removal of flatulence-causing factors from soymilk. *J. Microbiol. Biotechnol.* 20:1546-1554.
- Roy, I., A. Gupta, S. K. Khare, V. S. Bisaria and M. N. Gupta. 2003. Immobilization of xylan-degrading enzymes from *Melanocarpus albomyces* IIS 68 on the smart polymer Eudragit L-100. *Appl. Microbiol. Biotechnol.* 61:309-313.
- Shipkowski, S. and J. E. Brenchley. 2005. Characterization of an unusual cold-active beta-glucosidase belonging to family 3 of the glycoside hydrolases from the psychrophilic isolate *Paenibacillus* sp. strain C7. *Appl. Environ. Microbiol.* 71:4225-4232.
- Smiricky-Tjardes, M. R., C. M. Grieshop, E. A. Flickinger, L. L. Bauer and G. C. Fahey Jr. 2003. Dietary galactooligosaccharides affect ileal and total tract nutrient digestibility, ileal and fecal bacterial concentrations, and ileal fermentative characteristics of growing pigs. *J. Anim. Sci.* 81:2535-2545.
- Smits, C. H. M. and G. Annison. 1996. Non-starch plant polysaccharides in broiler nutrition-towards a physiologically valid approach to their determination. *World's Poult. Sci. J.* 52:204-221.
- Stein, H. H., L. L. Berger, J. K. Drackley, G. C. Fahey Jr, D. C. Hernot and C. M. Parsons. 2008. Nutrition properties and feeding values of soybeans and their coproducts. In: *Soybeans, Chemistry, Production, Processing, and Utilization* (Ed. A. Johnson, P. J. White and R. Galloway). AOCS press, Urbana, IL, USA. pp. 613-660.
- Syed, D. G., D. Agasar and A. Pandey. 2009. Production and partial purification of  $\alpha$ -amylase from a novel isolate *Streptomyces gulbargensis*. *J. Ind. Microbiol. Biotechnol.* 36:189-194.
- Tamura, K., J. Dudley, M. Nei and S. Kumar. 2007. MEGA4: Molecular evolution genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* 24:1596-1599.
- Tamura, K., M. Nei and S. Kumar. 2004. Prospect for inferring very large phylogenies by using the neighbor-joining methods. *Proc. Natl. Acad. Sci. USA.* 101:11030-11035.
- Thompson, J. D., D. G. Higgins and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22:4673-4680.
- Viana, P. A., S. T. De Rezende, V. M. Marques, L. M. Trevizano, F. M. L. Passos, M. G. A. Oliveira, M. P. Bemquerer, J. S. Oliveira and V. M. Guimaraes. 2006. Extracellular  $\alpha$ -galactosidase from *Debaryomyces hansenii* UFV-1 and its use in the hydrolysis of raffinose oligosaccharides. *J. Agric. Food Chem.* 54:2385-2391.
- Waeonukul, R., P. Pason, K. L. Kyu, K. Sakka, A. Kosugi, Y. Mori and K. Ratanakhanokchai. 2009. Cloning, sequencing, and expression of the gene encoding a multidomain endo-beta-1,4-xylanase from *Paenibacillus curdlanolyticus* B-6, and characterization of the recombinant enzyme. *J. Microbiol. Biotechnol.* 19:277-285.
- Wang, H., H. Luo, J. Li, Y. Bai, H. Huang, P. Shi, Y. Fan and B. Yao. 2010. An  $\alpha$ -galactosidase from an acidophilic *Bispora* sp. MEY-1 strain acts synergistically with  $\beta$ -mannase. *Bioresour. Technol.* 101:8376-8382.
- Wang, Q. F., J. L. Miao, Y. H. Hou, Y. Ding, G. D. Wang and G. Y. Li. 2005. Purification and characterization of an extracellular cold-active serine protease from the psychrophilic bacterium *Colwellia* sp. NJ341. *Biotechnol. Lett.* 27:1195-1198.
- Wang, Y., X. Gao, Q. Su, W. Wu and L. An. 2007. Cloning, expression, and enzyme characterization of an acid heat-stable phytase from *Aspergillus fumigatus* WY-2. *Curr. Microbiol.* 55:65-70.
- William, G. W., M. B. Susan, A. P. Dale and J. L. David. 1991. 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* 173:697-703.
- Yoon, M. Y. and H. J. Hwang. 2008. Reduction of soybean oligosaccharides and properties of  $\alpha$ -D-galactosidase from *Lactobacillus curvatus* R08 and *Leuconostoc mesenteriodes* JK55. *Food Microbiol.* 25:815-823.